

Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:sssptal644pnh  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Sep 29 The Philippines Inventory of Chemicals and Chemical  
Substances (PICCS) has been added to CHEMLIST  
NEWS 3 Oct 27 New Extraction Code PAX now available in Derwent  
Files  
NEWS 4 Oct 27 SET ABBREVIATIONS and SET PLURALS extended in  
Derwent World Patents Index files  
NEWS 5 Oct 27 Patent Assignee Code Dictionary now available  
in Derwent Patent Files  
NEWS 6 Oct 27 Plasdoc Key Serials Dictionary and Echoing added to  
Derwent Subscriber Files WPIDS and WPIX  
NEWS 7 Nov 29 Derwent announces further increase in updates for DWPI  
~~NEWS 8 Dec 5 French Multi-Disciplinary Database PASCAL Now on STN~~  
NEWS 9 Dec 5 Trademarks on STN - New DEMAS and EUMAS Files  
NEWS 10 Dec 15 2001 STN Pricing  
NEWS 11 Dec 17 Merged CEABA-VTB for chemical engineering and  
biotechnology  
NEWS 12 Dec 17 Corrosion Abstracts on STN  
NEWS 13 Dec 17 SYNTHLINE from Prous Science now available on STN  
NEWS 14 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 15 Jan 05 AIDSLINE is being removed from STN  
NEWS 16 Feb 06 Engineering Information Encompass files have new names  
  
NEWS EXPRESS FREE UPGRADE 5.0e FOR STN EXPRESS 5.0 WITH DISCOVER!  
(WINDOWS) NOW AVAILABLE  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 10:38:19 ON 12 FEB 2001 ✓

=> file medline embase biosis scisearch caplus

COST IN U.S. DOLLARS

SINCE FILE TOTAL  
ENTRY SESSION

Page 1

FULL ESTIMATED COST

0.15

0.15

FILE 'MEDLINE' ENTERED AT 10:38:50 ON 12 FEB 2001

FILE 'EMBASE' ENTERED AT 10:38:50 ON 12 FEB 2001

COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 10:38:50 ON 12 FEB 2001

COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'SCISEARCH' ENTERED AT 10:38:50 ON 12 FEB 2001

COPYRIGHT (C) 2001 Institute for Scientific Information (ISI) (R).

FILE 'CAPLUS' ENTERED AT 10:38:50 ON 12 FEB 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

=> s minor histocompatibility antigen

L1 2960 MINOR HISTOCOMPATIBILITY ANTIGEN

=> ~~s l1 and derivative~~

L2 19 L1 AND DERIVATIVE

=> dup remove 12

PROCESSING COMPLETED FOR L2

L3 13 DUP REMOVE L2 (6 DUPLICATES REMOVED)

=> d 13

L3 ANSWER 13 OF 13 MEDLINE

DUPLICATE 3

AN 86002336 MEDLINE

DN 86002336

TI **Minor histocompatibility antigens** are

developmentally regulated on murine embryonal carcinoma cells and their early differentiated **derivatives**.

AU Avner P; Simmler M C

SO CELL DIFFERENTIATION, (1985 Aug) 17 (2) 115-23.

Journal code: CQ6. ISSN: 0045-6039.

CY Ireland

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 198601

=> d l13 all 1-13

L13 NOT FOUND

The L-number entered has not been defined in this session, or it has been deleted. To see the L-numbers currently defined in this session, enter DISPLAY HISTORY at an arrow prompt (=>).

=> s 13

L4 13 L3

=> d 14 all 1113

13 ANSWERS ARE AVAILABLE. SPECIFIED ANSWER NUMBER EXCEEDS ANSWER SET SIZE

The answer numbers requested are not in the answer set.

ENTER ANSWER NUMBER OR RANGE (1):1-13

L4 ANSWER 1 OF 13 MEDLINE  
AN 2000269805 MEDLINE  
DN 20269805  
TI Adoptive immunotherapy in canine mixed chimeras after nonmyeloablative  
hematopoietic cell transplantation.  
AU Georges G E; Storb R; Thompson J D; Yu C; Gooley T; Bruno B; Nash R A  
CS Clinical Research Division, Fred Hutchinson Cancer Research Center,  
Department of Medicine, University of Washington, Seattle, WA 98109-1024,  
USA.. ggeorges@fhcrc.org  
NC DK42716 (NIDDK)  
CA15704 (NCI)  
CA78902 (NCI)

+  
SO BLOOD, (2000 May 15) 95 (10) 3262-9.  
Journal code: A8G. ISSN: 0006-4971.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
EM 200008  
EW 20000803  
AB Development of nontoxic and nonmyeloablative regimens for allogeneic  
hematopoietic stem-cell transplantation will decrease transplantation-  
related mortality caused by regimen-related toxic effects. In pursuit of  
this goal, a dog model of stable mixed hematopoietic chimerism was  
established in which leukocyte-antigen-identical litter mates are given  
sublethal total-body irradiation (2 Gy) before stem-cell transplantation  
and immunosuppression with mycophenolate mofetil and cyclosporine  
afterward. In the current study, we examined whether donor lymphocyte  
infusion (DLI) could be used as adoptive immunotherapy to convert mixed  
to complete donor chimerism. First, 8 mixed chimeras were given unmodified  
DLI between day 36 and day 414 after stem-cell transplantation. After a  
10- to 47-week follow-up period, there were no significant changes in the  
percentage of donor engraftment. Next, we immunized the donor to the  
**minor histocompatibility antigens** (mHA) of the  
recipient by means of repeated skin grafting. Lymphocytes from the  
mHA-sensitized donor were infused between day 201 and day 651 after  
transplantation. All 8 recipients of mHA-sensitized DLI had conversion to  
greater than 98% donor chimerism within 2 to 12 weeks of the infusion.  
Complications from mHA-sensitized DLI included graft-versus-host disease  
in 2 dogs and marrow aplasia in 1. These results showed that the low-dose  
transplant regimen establishes immune tolerance, and mHA-sensitized DLI  
is required to break tolerance, thereby converting mixed to complete donor  
chimerism. We propose that mixed chimerism established after  
nonmyeloablative allogeneic stem-cell transplantation provides a platform

of for adoptive immunotherapy that has clinical potential in the treatment  
 patients with malignant diseases.

CT Check Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.  
 Cyclosporine: AD, administration & dosage  
 Dogs  
 \*Hematopoietic Stem Cell Transplantation  
 Immunosuppressive Agents: AD, administration & dosage  
 \*Immunotherapy, Adoptive  
 Isoantigens  
 Lymphocyte Transfusion  
 \*Lymphocytes: IM, immunology  
**Mycophenolic Acid: AA, analogs & derivatives**  
 Mycophenolic Acid: AD, administration & dosage  
 Myeloablative Agonists: TU, therapeutic use  
 Tissue Donors  
 \*Transplantation Chimera  
 Transplantation Immunology

RN 128794-94-5 (RS 61443); 24280-93-1 (Mycophenolic Acid); 59865-13-3  
 (Cyclosporine)

CN 0 (Immunosuppressive Agents); 0 (Isoantigens); 0 (Myeloablative Agonists)

L4 ANSWER 2 OF 13 MEDLINE

AN 97256599 MEDLINE

DN 97256599

TI Identification of the rat maternally transmitted **minor**  
**histocompatibility antigen.**

AU Bhuyan P K; Young L L; Lindahl K F; Butcher G W

CS Howard Hughes Medical Institute, The University of Texas Southwestern  
 Medical Center, Dallas 75235, USA.

SO JOURNAL OF IMMUNOLOGY, (1997 Apr 15) 158 (8) 3753-60.

Journal code: IFB. ISSN: 0022-1767.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals

EM 199707

EW 19970702

AB The rat maternally transmitted Ag has been previously described as a  
 minor

histocompatibility Ag composed of a mitochondrially transmitted factor  
 (MTF) and the RT1.Aa MHC class I molecule. We compared the DNA sequences  
 of the 13 mitochondrial open reading frames from different rat strains

and

identified four coding polymorphisms that correlated with this MTF. We  
 used synthetic 17-mer peptides spanning the polymorphisms to sensitize  
 appropriate target cells in lymphocytotoxicity assays and found that the  
 MTF is derived from an internal region of ATPase 6. A tridecameric  
**derivative** of the ATPase 6 17 mer (termed 13N3E) could sensitize  
 RT1.Aa-expressing target cells at picomolar concentrations and, when  
 present on such cells, could compete fully with the natural ligand in  
 cold-target competition assays. Comparing the 13N3E peptide with the

known

peptide-binding requirements of RT1.Aa suggested two possible binding  
 conformations, placing either an internal or a C-terminal arginine in the  
 F pocket of the peptide-binding groove. Arguments favoring a "bulging"  
 conformation, with N- and C-terminal residues bound into their conserved  
 pockets, are discussed.

CT Check Tags: Animal; Female; Support, Non-U.S. Gov't  
 Amino Acid Sequence  
 DNA, Mitochondrial: GE, genetics  
 \*Immunity, Maternally-Acquired  
 \*Minor Histocompatibility Antigens: GE, genetics  
 \*Minor Histocompatibility Antigens: IM, immunology  
 Molecular Sequence Data  
 Pregnancy  
 Rats  
 Rats, Inbred Strains

CN 0 (DNA, Mitochondrial); 0 (Minor Histocompatibility Antigens)

L4 ANSWER 3 OF 13 MEDLINE  
 AN 96180356 MEDLINE  
 DN 96180356  
 TI Effect of metacycloprodigiosin, an inhibitor of killer T cells on murine skin and heart transplants.  
 AU Magae J; Miller M W; Nagai K; Shearer G M  
 CS Experimental Immunology Branch, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, USA.  
 SO JOURNAL OF ANTIBIOTICS, (1996 Jan) 49 (1) 86-90.  
~~Journal code: HCF. ISSN: 0021-8820.~~

---

CY Japan  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199608  
 AB Metacycloprodigiosin is an antibiotic that has been shown to suppress T-cell proliferation induced by concanavalin A in vitro. We examined the effect of metacycloprodigiosin on murine allogenic skin and heart transplantation models, and compared graft rejection with donor-specific cytotoxic T-cells and antibody activity. The antibiotic slightly prolonged the survival of C57Bl/6 heart and skin grafts in BALB/c mice, although the effect was less than that of cyclosporin A. The effect was more evident in Bm1 (H-2D mutant) skin grafts on C57Bl/6 hosts or in a minor histocompatibility antigen-mismatched model. In contrast, metacycloprodigiosin suppressed anti-graft cytotoxic T-cell activity of BALB/c spleen grafted with C57Bl/6 skin as comparable to cyclosporin A, but had only partial effect on antibody production. Thus, metacycloprodigiosin is more effective in reducing splenic cytotoxic T-cell activity than in prolonging murine skin or cardiac allografts.

CT Check Tags: Animal; Female  
 Cyclosporine: PD, pharmacology  
 Graft Survival: DE, drug effects  
 \*Heart Transplantation: IM, immunology  
 \*Immunosuppressive Agents: PD, pharmacology  
 Mice  
 Mice, Inbred BALB C  
 Mice, Inbred C57BL  
 \*Prodigiosin: AA, analogs & derivatives  
 Prodigiosin: PD, pharmacology  
 \*Skin Transplantation: IM, immunology  
 \*T-Lymphocytes, Cytotoxic: DE, drug effects  
 T-Lymphocytes, Cytotoxic: IM, immunology

RN 59865-13-3 (Cyclosporine); 82-89-3 (Prodigiosin)  
 CN 0 (metacycloprodigiosin); 0 (Immunosuppressive Agents)

L4 ANSWER 4 OF 13 MEDLINE  
 AN 95002951 MEDLINE  
 DN 95002951  
 TI Inhibition of nitric oxide production is associated with enhanced weight loss, decreased survival, and impaired alloengraftment in mice undergoing graft-versus-host disease after bone marrow transplantation.  
 AU Drobyski W R; Keever C A; Hanson G A; McAuliffe T; Griffith O W  
 CS Department of Medicine, Medical College of Wisconsin, Milwaukee 53226.  
 NC CA01534 (NCI)  
 SO BLOOD, (1994 Oct 1) 84 (7) 2363-73.  
 Journal code: A8G. ISSN: 0006-4971.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
 EM 199501  
 AB The pathophysiologic role of nitric oxide (NO) in graft-versus-host disease (GVHD) was investigated in a murine bone marrow (BM) transplantation model where donor and recipient were H-2-matched but differed at multiple ~~minor histocompatibility~~  
**antigens.** Host AKR/J (H-2K) mice received lethal total body irradiation as pretransplant conditioning followed by transplantation of donor B10.BR (H-2K) BM cells with or without spleen cells as a source of GVH-reactive T cells. NO production, as assessed by serum nitrate and nitrite levels, was increased for up to 3 weeks posttransplant in animals undergoing both moderate and severe GVHD. Administration of NG-methyl-L-arginine (L-NMA), an inhibitor of nitric oxide synthase, to animals undergoing GVHD resulted in effective suppression of NO  
 production  
 when compared with saline-treated GVHD control animals. Suppression of NO production by L-NMA in GVHD animals was associated with enhanced weight loss early posttransplant and decreased overall survival. Histologic analysis of tissues from L-NMA-treated and saline-treated GVHD animals showed that early weight loss was not because of an exacerbation of GVHD, indicating that NO did not appear to play an immunosuppressive role in this experimental model. L-NMA-treated animals with enhanced weight loss were observed to have splenic atrophy, decreased extramedullary hematopoiesis, and a reduction in BM cellularity when compared with GVHD control mice that were weight-matched before transplant. Analysis of T-cell chimerism in the spleen showed that L-NMA treatment impaired donor T-cell repopulation. In vitro colony-forming unit (CFU) assays were performed to further assess the role of NO on BM progenitor cell growth. L-NMA added directly into culture had no effect on CFU-granulocyte/macrophage (CFU-GM) formation in normal murine BM. In contrast, total CFU-GM from L-NMA-treated animals were significantly reduced when compared with GVHD controls or BM control animals who did  
 not  
 develop GVHD. Collectively, these data indicate that inhibition of NO impairs hematopoietic reconstitution and support the premise that NO appears to play a novel role in the facilitation of alloengraftment posttransplant.  
 CT Check Tags: Animal; Support, U.S. Gov't, P.H.S.  
 Amino Acid Oxidoreductases: AI, antagonists & inhibitors  
**Arginine: AA, analogs & derivatives**  
 Arginine: PD, pharmacology

Body Weight: DE, drug effects  
 Bone Marrow: PA, pathology  
 \*Bone Marrow Transplantation: PA, pathology  
 \*Graft vs Host Disease: PA, pathology  
 Graft Survival  
 Mice  
 Mice, Inbred AKR  
 Minor Lymphocyte Stimulatory Antigens: IM, immunology  
 \*Nitric Oxide: BI, biosynthesis  
 Spleen: PA, pathology  
 Survival Analysis  
 RN 10102-43-9 (Nitric Oxide); 17035-90-4 (omega-N-Methylarginine); 7004-12-8 (Arginine)  
 CN EC 1.14.13.39 (Nitric-Oxide Synthase); EC 1.4. (Amino Acid Oxidoreductases); 0 (Minor Lymphocyte Stimulatory Antigens)  
 L4 ANSWER 5 OF 13 MEDLINE  
 AN 86002336 MEDLINE  
 DN 86002336  
 TI **Minor histocompatibility antigens are developmentally regulated on murine embryonal carcinoma cells and their early differentiated derivatives.**  
 AU ~~Avner P; Simmler M C~~  
 SO CELL DIFFERENTIATION, (1985 Aug) 17 (2) 115-23.  
 Journal code: CQ6. ISSN: 0045-6039.  
 CY Ireland  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 198601  
 AB Differences in the expression of minor histocompatibility (Hm) alloantigens on two mouse embryonal carcinoma (EC) cell lines and the PYS-2 and T.D.M.-1 differentiated **derivatives** have been demonstrated by their ability to elicit a cytolytic T lymphocyte response.  
 Experiments involving the use of various responder-target strain combinations and recombinant inbred mice strains have shown that: (1) there are major differences in Hm expression on EC cells compared with differentiated **derivatives** whose Hm expression appears more like that of adult splenocytes; (2) although both EC cell lines show reduced Hm immunogenicity compared with adult splenocytes, major differences in the expression and possible presentation of Hm between the F9 and PCC3 EC cell lines can be detected by in vivo priming and by in vitro cold competition target experiments. These observations are discussed in relation to the differences in allograft rejection patterns observed with PCC3 and F9 and to possible differences in developmental staging of these cell lines.  
 CT Check Tags: Animal  
 Cell Differentiation  
 Cell Line  
 Cytotoxicity, Immunologic  
 Immunization  
 Mice  
 Mice, Inbred Strains  
 \*Minor Histocompatibility Loci  
 T-Lymphocytes, Cytotoxic: IM, immunology  
 \*Teratoma: IM, immunology

Teratoma: PA, pathology  
Yolk Sac: CY, cytology  
Yolk Sac: IM, immunology

L4 ANSWER 6 OF 13 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
AN 97330017 EMBASE  
DN 1997330017  
TI Inducible nitric oxide synthase suppresses the development of allograft arteriosclerosis.  
AU Shears II L.L.; Kawaharada N.; Tzeng E.; Billiar T.R.; Watkins S.C.; Kovesdi I.; Lizonova A.; Pham S.M.  
CS Dr. S.M. Pham, Presbyterian University Hospital, 200 Lothrop Street, Pittsburgh, PA 15213, United States. pham@pittsburg.nb.upmc.edu  
SO Journal of Clinical Investigation, (1997) 100/8 (2035-2042).  
Refs: 48  
ISSN: 0021-9738 CODEN: JCINAO  
CY United States  
DT Journal; Article  
FS 018 Cardiovascular Diseases and Cardiovascular Surgery  
037 Drug Literature Index  
LA English  
SL English  
AB ~~In cardiac transplantation~~, chronic rejection takes the form of an occlusive vasculopathy. The mechanism underlying this disorder remains unclear. The purpose of this study was to investigate the role nitric oxide (NO) may play in the development of allograft arteriosclerosis. Rat aortic allografts from ACI donors to Wistar Furth recipients with a strong genetic disparity in both major and minor histocompatibility antigens were used for transplantation. Allografts collected at 28 d were found to have significant increases in both inducible NO synthase (iNOS) mRNA and protein as well as in intimal thickness when compared with isografts. Inhibiting NO production with an iNOS inhibitor increased the intimal thickening by 57.2%, indicating that NO suppresses the development of allograft arteriosclerosis. Next, we evaluated the effect of cyclosporine (CsA) on iNOS expression and allograft arteriosclerosis. CsA (10 mg/kg/d) suppressed the expression of iNOS in response to balloon-induced aortic injury. Similarly, CsA inhibited iNOS expression in the aortic allografts, associated with a 65% increase in intimal thickening. Finally, we investigated the effect of adenoviral-mediated iNOS gene transfer on allograft arteriosclerosis. Transduction with iNOS using an adenoviral vector suppressed completely the development of allograft arteriosclerosis in both untreated recipients and recipients treated with CsA. These results suggest that the early immune-mediated upregulation in iNOS expression partially protects aortic allografts from the development of allograft arteriosclerosis, and that iNOS gene transfer strategies may prove useful in preventing the development of this otherwise untreatable disease process.  
CT Medical Descriptors:  
\*atherosclerosis: CO, complication  
\*atherosclerosis: PC, prevention  
\*atherosclerosis: ET, etiology  
\*graft rejection: PC, prevention  
\*graft rejection: ET, etiology  
\*graft rejection: DT, drug therapy



\*graft rejection: CO, complication  
 \*heart transplantation  
 adenovirus  
 allograft  
 animal model  
 animal tissue  
 artery intima proliferation: ET, etiology  
 artery intima proliferation: PC, prevention  
 artery intima proliferation: CO, complication  
 article  
 controlled study  
 enzyme induction  
 gene transfer  
 graft failure  
 immunosuppressive treatment  
 nonhuman  
 priority journal  
 rat  
 subcutaneous drug administration  
 virus vector  
 Drug Descriptors:  
 \*cyclosporin a: DT, drug therapy  
~~\*nitric oxide: EC, endogenous compound~~  
 \*nitric oxide synthase: EC, endogenous compound  
 immunosuppressive agent: DT, drug therapy  
**lysine derivative**  
 messenger rna: EC, endogenous compound

RN (cyclosporin a) 59865-13-3, 63798-73-2; (nitric oxide) 10102-43-9;  
 (nitric  
 oxide synthase) 125978-95-2

L4 ANSWER 7 OF 13 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
 AN 94028927 EMBASE  
 DN 1994028927  
 TI Veto suppression: The peripheral way of T cell tolerization.  
 AU Tscherning T.; Claesson M.H.  
 CS Lab of Experimental Immunology, Institute of Medical Anatomy, The Panum  
 Institute, Blegdamsvej 3C, DK-2200 Copenhagen N, Denmark  
 SO Experimental and Clinical Immunogenetics, (1993) 10/4 (179-188).  
 ISSN: 0254-9670 CODEN: ECIME4  
 CY Switzerland  
 DT Journal; General Review  
 FS 022 Human Genetics  
 026 Immunology, Serology and Transplantation  
 LA English  
 SL English  
 AB Cells with veto activity induce a state of tolerance in T cell precursors  
 with specificity for antigen determinants expressed on the surface of the  
 veto-active cell. This state of tolerance is not strictly defined, but  
 results in altered responses to specific antigen, such as decreased  
 proliferation, decreased development of cytotoxicity and secretion of  
 interleukins, down-regulated ability to reject grafts and expression of T  
 cell and IL-2 receptors. Both clonal anergy and clonal deletion has been  
 shown to operate in vetoed T cells. Veto-induced tolerance can be  
 established in vitro and in vivo for both MHC class I and II as well as  
**minor histocompatibility antigens**. The most  
 powerful veto activity is present in mature activated cytotoxic CD8+ T  
 cells, but other cells including noncytotoxic cells are also capable of

acting as veto cells. Thus it appears that veto activity per se is not confined to a certain cellular entity, but rather reflects a constitutively expressed immunoregulatory capability inherent to a broad array of activated T cell and non-T cell categories with their own distinct functions not related to their eventual veto activity.

CT Medical Descriptors:

\*immunological tolerance

\*immunoregulation

\*t lymphocyte

clonal anergy

cell proliferation

cytotoxicity

graft rejection

lymphocyte clone

mouse

nonhuman

precursor cell

priority journal

protein secretion

review

Drug Descriptors:

t lymphocyte receptor

cd8 antigen: EC, endogenous compound

interleukin 2 receptor: EC, endogenous compound

**interleukin derivative: EC, endogenous compound**

major histocompatibility antigen class 1: EC, endogenous compound

major histocompatibility antigen class 2: EC, endogenous compound

membrane antigen: EC, endogenous compound

L4 ANSWER 8 OF 13 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1990:28060 BIOSIS

DN BA89:15026

TI PROTEIN-SPECIFIC CYTOTOXIC T LYMPHOCYTES RECOGNITION OF TRANSFECTANTS EXPRESSING INTRACELLULAR MEMBRANE-ASSOCIATED OR SECRETED FORMS OF BETA GALACTOSIDASE.

AU RAMMENSEE H-G; SCHILD H; THEOPOLD U

CS MAX-PLANCK-INST. BIOL., ABT. IMMUNGENETIK, CORRENSSTRASSE 42, D-7400 TUEBINGEN, FRG.

SO IMMUNOGENETICS, (1989) 30 (4), 296-302.

CODEN: IMNGBK. ISSN: 0093-7711.

FS BA; OLD

LA English

AB BALB/c-derived tumor cells were transfected with recombinant Escherichia coli .beta.-galactosidase (.beta.-gal) gene which were inserted into IgM heavy chain gene **derivatives**, leading to expression of the resulting fusion protein in different cellular compartments. A .beta.-gal-specific, major histocompatibility complex (MHC) class I-restricted CD8+ CD4- cytotoxic T lymphocyte (CTL) line of BALB/c origin raised against one transfectant expressing cytoplasmic .beta.-gal also lysed transfectants expressing .beta.-gal as membrane-inserted fusion protein, as well as transfectants secreting .beta.-gal. Our data show

that

MHC class I-restricted CTL can recognize fragments of nonviral cellular proteins, be they expressed as intracellular, membrane-inserted, or secreted products. The findings confirm and extend a hypothesis on the nature of minor histocompatibility (H) antigens formulated earlier.

CC Cytology and Cytochemistry - Animal \*02506

Biochemical Studies - Proteins, Peptides and Amino Acids 10064

Biophysics - Membrane Phenomena \*10508  
Enzymes - Physiological Studies \*10808  
Metabolism - Proteins, Peptides and Amino Acids \*13012  
Immunology and Immunochemistry - Immunopathology, Tissue Immunology  
\*34508

BC Muridae 86375

IT Miscellaneous Descriptors

MOUSE MAJOR HISTOCOMPATIBILITY COMPLEX MINOR

**HISTOCOMPATIBILITY ANTIGENS**

RN 9031-11-2 (BETA GALACTOSIDASE)

L4 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2001 ACS

AN 1999:96271 CAPLUS

DN 130:167164

TI The HA-1 antigen

IN Goulmy, Elsa Afra Julia Maria; Hunt, Donald F.; Engelhard, Victor H.

PA Rijksuniversiteit te Leiden, Neth.

SO PCT Int. Appl., 47 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07K014-705

~~ICS C07K016-28; A61K038-17~~

CC 15-2 (Immunochemistry)

FAN.CNT 3

|      | PATENT NO.     | KIND   | DATE     | APPLICATION NO. | DATE     |
|------|----------------|--|----------|-----------------|----------|
| PI   | WO 9905174     | A1   | 19990204 | WO 1998-NL425   | 19980723 |
|      | W:             | AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |          |                 |          |
|      | RW:            | GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG   |          |                 |          |
|      | AU 9885640     | A1   | 19990216 | AU 1998-85640   | 19980723 |
| PRAI | EP 1997-202303 |  | 19970723 |                 |          |
|      | WO 1998-N      |  |          |                 |          |

L425 19980723

AB The present invention discloses the peptide sequence of a so called minor H antigen. The minor H antigens are assocd. with the graft vs. host disease. The peptide and its **derivs.** find many uses in bone marrow transplantation, organ transplantation and in the treatment of leukemia. The peptide and its **derivs.** can be incorporated in vaccines, in pharmaceutical formulations and they can be used in diagnostic test kits. The peptide is derived from the HA-1 minor antigen and has the sequence VLXDDLLEA, wherein X represents a histidine or an arginine residue. Both donors and recipients in bone marrow transplantation can be treated with the peptides, optionally in combination with other peptides, coupled to carriers, with suitable excipients and/or adjuvants.

ST **minor histocompatibility antigen** HA1 immune

tolerance; T cell epitope HA1 antigen leukemia; graft vs host disease

IT **Minor histocompatibility antigens**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(HA-1; T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Anti-idiotypic antibodies  
 Autoimmune diseases  
 B cell (lymphocyte)  
 Bone marrow transplant  
 Drug delivery systems  
 Graft vs. host reaction  
 Immune tolerance  
 Immunization  
 Immunological diseases  
 Leukemia  
 Mammal (Mammalia)  
 Medicine  
 Protein sequences  
 Transplant (organ)  
 Transplant rejection  
 Vaccines  
 (T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

---

IT Antibodies  
 TCR (T cell receptors)  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL  
 (Biological study); PREP (Preparation); USES (Uses)  
 (T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Class I HLA antigens  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Epitopes  
 (T cell; T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Test kits  
 (diagnostic; T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT T cell (lymphocyte)  
 (epitope; T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Hematopoietic precursor cell  
 (tumors; T cell epitope obtainable from the **minor**  
**histocompatibility antigen HA-1** for induction of  
 immune tolerance and for treating transplant rejection, autoimmune  
 disease, neoplastic hematopoietic disease, and graft vs host disease)

IT 204931-32-8    220419-68-1

RL: BSU (Biological study, unclassified); PRP (Properties); THU  
(Therapeutic use); BIOL (Biological study); USES (Uses)  
(T cell epitope obtainable from the **minor**  
**histocompatibility antigen** HA-1 for induction of  
immune tolerance and for treating transplant rejection, autoimmune  
disease, neoplastic hematopoietic disease, and graft vs host disease)

RE.CNT 6

RE

- (1) Den Haan, J; Eur J Immunol 1996, V26, P2680 CAPLUS
- (2) Den Haan, J; Science 1995, V268, P1476 CAPLUS
- (3) Den Haan, J; Science 1998, V279, P1054112
- (4) Goulmy, E; WO 9705168 A 1997 CAPLUS
- (5) Goulmy, E; WO 9705169 A 1997 CAPLUS
- (6) Goulmy, E; Eye 1995, V9, P180

L4 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2001 ACS

AN 1999:96270 CAPLUS

DN 130:167163

TI The HA-1 antigen

IN Goulmy, Elsa Afra Julia Maria; Hunt, Donald Frederick; Engelhard, Victor Henry

PA Rijksuniversiteit te Leiden, Neth.

SO PCT Int. Appl., 57 pp

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07K014-705

ICS C07K016-28; A61K038-17; C12N005-06

CC 15-2 (Immunochemistry)

Section cross-reference(s): 3

FAN.CNT 3

|    | PATENT NO. | KIND   | DATE     | APPLICATION NO. | DATE     |
|----|------------|--|----------|-----------------|----------|
| PI | WO 9905173 | A1   | 19990204 | WO 1998-NL424   | 19980723 |
|    | W:         | AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |          |                 |          |
|    | RW:        | GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG   |          |                 |          |
|    | AU 9885639 | A1   | 19990216 | AU 1998-85639   | 19980723 |
|    | EP 996636  | A1   | 20000503 | EP 1998-936758  | 19980723 |
|    | R:         | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  |          |                 |          |

IE

PRAI EP 1997-202303 19970723

WO 1998-N

L424 19980723

AB The present invention discloses the peptide sequence of a so-called minor H antigen. The minor H antigens are assocd. with the graft vs. host disease. The peptide and its **derivs.** find many uses in bone marrow transplantation, organ transplantation and in the treatment of leukemia. The peptide and its **derivs.** can be incorporated in vaccines, in pharmaceutical formulations and they can be used in diagnostic test kits. The peptide is derived from the HA-1 minor antigen and has the sequence VLXDDLLEA, wherein X represents a histidine or an arginine residue. Both donors and recipients in bone marrow

transplantation can be treated with the peptides, optionally in combination with other peptides, coupled to carriers, with suitable excipients and/or adjuvants.

ST **minor histocompatibility antigen** HA1 immune tolerance; T cell epitope HA1 transplant rejection; graft vs host disease HA1 antigen

IT **Minor histocompatibility antigens**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(HA-1; T cell epitope of **minor histocompatibility**

**antigen** HA-1 for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia

and

immune disease)

IT Genes (animal)

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(KIAA0223; T cell epitope of **minor histocompatibility**

**antigen** HA-1 for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia

and

immune disease)

IT Anti-idiotypic antibodies

~~Autoimmune diseases~~

B cell (lymphocyte)

Bone marrow transplant

Cytotoxic T cell

Dendritic cell

Drug delivery systems

Epitopes

Graft vs. host reaction

Hematopoietic precursor cell

Immune tolerance

Immunization

Immunological diseases

Mammal (Mammalia)

Medicine

Polymorphism (genetic)

T cell (lymphocyte)

Transplant (organ)

Transplant rejection

Vaccines

(T cell epitope of **minor histocompatibility**

**antigen** HA-1 for induction of immune tolerance and for

treatment of transplant rejection, graft vs. host disease, leukemia

and

immune disease)

IT Antibodies

TCR (T cell receptors)

RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL

(Biological study); PREP (Preparation); USES (Uses)

(T cell epitope of **minor histocompatibility**

**antigen** HA-1 for induction of immune tolerance and for

treatment of transplant rejection, graft vs. host disease, leukemia

and

immune disease)

IT Class I HLA antigens

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(T cell epitope of **minor histocompatibility**

antigen HA-1 for induction of immune tolerance and for  
treatment of transplant rejection, graft vs. host disease, leukemia  
and  
immune disease)  
IT Genes (animal)  
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL  
(Biological study); USES (Uses)  
(suicide; T cell epitope of **minor histocompatibility**  
**antigen** HA-1 for induction of immune tolerance and for  
treatment of transplant rejection, graft vs. host disease, leukemia  
and  
immune disease)  
IT Hematopoietic precursor cell  
(tumors; T cell epitope of **minor histocompatibility**  
**antigen** HA-1 for induction of immune tolerance and for  
treatment of transplant rejection, graft vs. host disease, leukemia  
and  
immune disease)  
IT 204931-32-8 220419-68-1  
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study);  
USES  
(Uses)

~~(T cell epitope of **minor histocompatibility**~~

**antigen** HA-1 for induction of immune tolerance and for  
treatment of transplant rejection, graft vs. host disease, leukemia  
and  
immune disease)

RE.CNT 7

RE

- (1) Den Haan, J; Eur J Immunol 1996, V26, P2680 CAPLUS
- (2) Den Haan, J; Science 1995, V268, P1476 CAPLUS
- (3) Den Haan, J; Science 1998, V279, P1054 CAPLUS
- (4) Goulmy, E; WO 9705168 A 1997 CAPLUS
- (5) Goulmy, E; WO 9705169 A 1997 CAPLUS
- (6) Goulmy, E; Eye 1995, V9, P180
- (7) Van Der Harst, D; Blood 1994, V83(4), P1060 CAPLUS

L4 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2001 ACS

AN 1997:215796 CAPLUS

DN 126:198552

TI HA-2 antigenic peptide

IN Goulmy, Els A. J. M.; Hunt, Donald F.; Engelhard, Victor H.

PA Rijksuniversiteit Te Leiden, Neth.; University of Virginia Patent  
Foundation; Goulmy, Els A. J. M.; Hunt, Donald F.; Engelhard, Victor H.

SO PCT Int. Appl., 36 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07K014-74

ICS C07K016-28; A61K038-16; C12N005-08

CC 15-2 (Immunochemistry)

FAN.CNT 1

|    | PATENT NO. | KIND  | DATE     | APPLICATION NO. | DATE     |
|----|------------|---|----------|-----------------|----------|
| PI | WO 9705169 | A1  | 19970213 | WO 1996-NL183   | 19960425 |
|    | W:         | AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, |          |                 |          |

SG, SI  
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR,  
IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN

|            |    |          |                 |          |
|------------|----|----------|-----------------|----------|
| US 5770201 | A  | 19980623 | US 1994-363691  | 19941223 |
| CA 2224909 | AA | 19970213 | CA 1996-2224909 | 19960425 |
| AU 9654099 | A1 | 19970226 | AU 1996-54099   | 19960425 |
| AU 716907  | B2 | 20000309 |                 |          |
| EP 840750  | A1 | 19980513 | EP 1996-911119  | 19960425 |

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
IE, SI, LT, LV, FI

|             |    |          |                |          |
|-------------|----|----------|----------------|----------|
| JP 11514340 | T2 | 19991207 | JP 1996-507492 | 19960425 |
|-------------|----|----------|----------------|----------|

PRAI EP 1995-202039 19950725  
WO 1996-N  
L183 19960425

AB The present invention discloses the first peptide sequence of a so-called  
minor H antigen. The minor H antigens are assocd. with the Graft vs.  
Host

Disease. The peptide and its **derivs.** find many uses in bone  
marrow transplantation, organ transplantation and in the treatment of  
leukemia. The peptide and its **derivs.** can be incorporated in  
vaccines, in pharmaceutical formulations and they can be used in  
diagnostic test kits. The peptide is derived from the HA-2 minor antigen  
~~and has the sequence IXGEVXVSV, wherein X represents a leucine or an~~  
isoleucine residue. Both donors and recipients in bone marrow  
transplantation can be treated with the peptides, optionally in  
combination with other peptides, coupled to carriers, with suitable  
excipients and/or adjuvants.

ST **minor histocompatibility antigen HA2**  
transplant rejection

IT **Minor histocompatibility antigens**  
RL: BSU (Biological study, unclassified); PRP (Properties); THU  
(Therapeutic use); BIOL (Biological study); USES (Uses)  
(HA-2; **minor histocompatibility antigen**  
HA-2 peptide for treating transplant rejection)

IT T cell (lymphocyte)  
(epitope; **minor histocompatibility antigen**  
HA-2 peptide for treating transplant rejection)

IT B cell (lymphocyte)  
Graft-vs.-host reaction  
Immune tolerance  
Leukemia  
Protein sequences  
Transplant rejection  
Vaccines  
(**minor histocompatibility antigen HA-2**  
peptide for treating transplant rejection)

IT TCR (T-cell receptors)  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(**minor histocompatibility antigen HA-2**  
peptide for treating transplant rejection)

IT Antibodies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(**minor histocompatibility antigen HA-2**  
peptide for treating transplant rejection)

IT Hematopoietic precursor cell  
(tumors; **minor histocompatibility antigen**  
HA-2 peptide for treating transplant rejection)

IT 187944-95-2 187944-96-3 187944-97-4



RL: BSU (Biological study, unclassified); PRP (Properties); THU  
(Therapeutic use); BIOL (Biological study); USES (Uses)  
(**minor histocompatibility antigen HA-2**  
peptide for treating transplant rejection)

L4 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2001 ACS  
AN 1990:1427 CAPLUS  
DN 112:1427  
TI Mapping minor H genes  
AU Simpson, E.; Tomonari, K.  
CS Transplant. Biol. Sect., MRC Clin. Res. Cent., Harrow/Middlesex, HA1 3UJ,  
UK  
SO Immunology (1989), Suppl. 2, 42-9  
CODEN: IMMUAM; ISSN: 0019-2805  
DT Journal; General Review  
LA English  
CC 3-0 (Biochemical Genetics)  
Section cross-reference(s): 13, 15  
AB A review with 14 refs. The manner in which minor histocompatibility (H)  
antigens have been defined in mouse and man, in vivo and in vitro, is  
considered. Chromosomal mapping of minor H genes using T-cell clones is  
illustrated, with particular ref. to the H-Y antigen gene, using the  
~~sex-reversing translocation Sxr of mouse and the deriv. Sxr'~~  
mutation. A no. of minor H antigen-specific T-cell clones restricted by  
class I or class II major histocompatibility complex (MHC) mols. are  
described, together with information about their phenotypes and T-cell  
receptor usage.  
ST histocompatibility H antigen gene mapping review  
IT Gene and Genetic element, animal  
RL: BIOL (Biological study)  
(for **minor histocompatibility antigens**,  
mapping of)  
IT Antigens  
RL: BIOL (Biological study)  
(H, genes for minor, mapping of)

L4 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2001 ACS  
AN 1985:521308 CAPLUS  
DN 103:121308  
TI Differential Hm antigen expression on EC cells and early differentiated  
**derivatives**  
AU Simmler, M. C.; Avner, P. R.  
CS Unite Immunol. Virol. Tumeurs, Hop. Cochin, Paris, 75014, Fr.  
SO EMBO J. (1985), 4(5), 1177-85  
CODEN: EMJODG; ISSN: 0261-4189  
DT Journal  
LA English  
CC 15-2 (Immunochemistry)  
AB Differences in the expression of minor histocompatibility (Hm)  
alloantigens on 2 mouse embryonal carcinoma (EC) cell lines and the PYS-2  
and T.D.M.-1 differentiated **derivs.** have been demonstrated by  
their ability to elicit a cytolytic T-lymphocyte (CTL) response. Expts.  
involving the use of various responder-target strain combinations on the  
one hand and recombinant inbred (RI) mice strains on the other have shown  
that: (i) there are major differences in Hm expression on the EC cells  
compared with the differentiated **derivs.** whose Hm expression  
appears more akin to that of adult splenocytes; (ii) although both EC  
cell

lines show reduced Hm immunogenicity compared with adult splenocytes, major differences in the expression and possibly presentation between the F9 and PCC3 EC cell lines can be detected both by in vivo priming and by in vitro cold competition target expts. These results are discussed in connection with the unexpected finding that some EC cell lines are capable of specific competition effects for appropriate CTL effectors despite their inability to stimulate such effectors in vitro and the absence of major histocompatibility complex products.

ST **minor histocompatibility antigen** embryonal carcinoma

IT Lymphocyte  
(T-, cytolytic, **minor histocompatibility antigen** expression on embryonal carcinoma cells in relation to)

IT Carcinoma  
(embryonal, **minor histocompatibility antigen** expression on, cytolytic T lymphocyte response in relation to)

IT Antigens  
RL: BIOL (Biological study).  
(minor histocompatibility, of embryonal carcinoma cells, cytolytic T lymphocyte response in relation to)

---

=> d his

(FILE 'HOME' ENTERED AT 10:38:19 ON 12 FEB 2001)

FILE 'MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS' ENTERED AT 10:38:50 ON 12 FEB 2001

L1 2960 S MINOR HISTOCOMPATIBILITY ANTIGEN  
L2 19 S L1 AND DERIVATIVE  
L3 13 DUP REMOVE L2 (6 DUPLICATES REMOVED)  
L4 13 S L3

=> s l1 and HA-1

L5 149 L1 AND HA-1

=> dup remove l5

PROCESSING COMPLETED FOR L5

L6 58 DUP REMOVE L5 (91 DUPLICATES REMOVED)

=> s l6 and VLXDDLLEA

L7 2 L6 AND VLXDDLLEA

=> d l7 all 1-2

L7 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2001 ACS  
AN 1999:96271 CAPLUS  
DN 130:167164  
TI The **HA-1** antigen  
IN Goulmy, Elsa Afra Julia Maria; Hunt, Donald F.; Engelhard, Victor H.  
PA Rijksuniversiteit te Leiden, Neth.

SO PCT Int. Appl., 47 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07K014-705  
 ICS C07K016-28; A61K038-17  
 CC 15-2 (Immunochemistry)  
 FAN.CNT 3

|      | PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|------|---|------|----------|-----------------|----------|
| PI   | WO 9905174  | A1   | 19990204 | WO 1998-NL425   | 19980723 |
|      | W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE,<br>DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG,<br>KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX,<br>NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,<br>UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM<br>RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES,<br>FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI,<br>CM, GA, GN, GW, ML, MR, NE, SN, TD, TG |      |          |                 |          |
|      | AU 9885640  | A1   | 19990216 | AU 1998-85640   | 19980723 |
| PRAI | EP 1997-202303  |      | 19970723 |                 |          |
|      | WO 1998-N   |      |          |                 |          |
| I425 | 19980723  |      |          |                 |          |

AB The present invention discloses the peptide sequence of a so called minor H antigen. The minor H antigens are assocd. with the graft vs. host disease. The peptide and its derivs. find many uses in bone marrow transplantation, organ transplantation and in the treatment of leukemia. The peptide and its derivs. can be incorporated in vaccines, in pharmaceutical formulations and they can be used in diagnostic test kits. The peptide is derived from the HA-1 minor antigen and has the sequence VLXDDLLEA, wherein X represents a histidine or an arginine residue. Both donors and recipients in bone marrow transplantation can be treated with the peptides, optionally in combination with other peptides, coupled to carriers, with suitable excipients and/or adjuvants.

ST **minor histocompatibility antigen** HA1 immune tolerance; T cell epitope HA1 antigen leukemia; graft vs host disease

IT **Minor histocompatibility antigens**  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (HA-1; T cell epitope obtainable from the **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Anti-idiotypic antibodies  
 Autoimmune diseases  
 B cell (lymphocyte)  
 Bone marrow transplant  
 Drug delivery systems  
 Graft vs. host reaction  
 Immune tolerance  
 Immunization  
 Immunological diseases  
 Leukemia  
 Mammal (Mammalia)  
 Medicine  
 Protein sequences

Transplant (organ)  
Transplant rejection  
Vaccines

(T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Antibodies

TCR (T cell receptors)

RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Class I HLA antigens

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Epitopes

(T cell; T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Test kits

(diagnostic; T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT T cell (lymphocyte)

(epitope; T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT Hematopoietic precursor cell:

(tumors; T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

IT 204931-32-8 220419-68-1

RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(T cell epitope obtainable from the **minor**

**histocompatibility antigen HA-1**

for induction of immune tolerance and for treating transplant rejection, autoimmune disease, neoplastic hematopoietic disease, and graft vs host disease)

RE.CNT 6

RE

- (1) Den Haan, J; Eur J Immunol 1996, V26, P2680 CAPLUS
- (2) Den Haan, J; Science 1995, V268, P1476 CAPLUS
- (3) Den Haan, J; Science 1998, V279, P1054112
- (4) Goulmy, E; WO 9705168 A 1997 CAPLUS
- (5) Goulmy, E; WO 9705169 A 1997 CAPLUS
- (6) Goulmy, E; Eye 1995, V9, P180

L7 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2001 ACS  
 AN 1999:96270 CAPLUS  
 DN 130:167163  
 TI The HA-1 antigen  
 IN Goulmy, Elsa Afra Julia Maria; Hunt, Donald Frederick; Engelhard, Victor Henry  
 PA Rijksuniversiteit te Leiden, Neth.  
 SO PCT Int. Appl., 57 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07K014-705  
 ICS C07K016-28; A61K038-17; C12N005-06  
 CC 15-2 (Immunochemistry)  
 Section cross-reference(s): 3  
 FAN.CNT 3

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| PI WO 9905173   | A1   | 19990204 | WO 1998-NL424   | 19980723 |
| W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM<br>RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG |      |          |                 |          |
| AU 9885639  | A1   | 19990216 | AU 1998-85639   | 19980723 |
| EP 996636   | A1   | 20000503 | EP 1998-936758  | 19980723 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  |      |          |                 |          |

IE

PRAI EP 1997-202303 19970723

WO 1998-N

L424 19980723

AB The present invention discloses the peptide sequence of a so-called minor H antigen. The minor H antigens are assocd. with the graft vs. host disease. The peptide and its derivs. find many uses in bone marrow transplantation, organ transplantation and in the treatment of leukemia. The peptide and its derivs. can be incorporated in vaccines, in pharmaceutical formulations and they can be used in diagnostic test kits. The peptide is derived from the HA-1 minor antigen and has the sequence VLXDDLLEA, wherein X represents a histidine or an arginine residue. Both donors and recipients in bone marrow transplantation can be treated with the peptides, optionally in combination with other peptides, coupled to carriers, with suitable excipients and/or adjuvants.

ST **minor histocompatibility antigen** HA1 immune tolerance; T cell epitope HA1 transplant rejection; graft vs host disease HA1 antigen

IT **Minor histocompatibility antigens**

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL

(Biological study); USES (Uses)  
 (HA-1; T cell epitope of **minor histocompatibility antigen HA-1**  
 for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Genes (animal)  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (KIAA0223; T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Anti-idiotypic antibodies  
 Autoimmune diseases  
 B cell (lymphocyte)  
 Bone marrow transplant  
 Cytotoxic T cell  
 Dendritic cell  
 Drug delivery systems  
 Epitopes  
 Graft vs. host reaction  
 Hematopoietic precursor cell  
 Immune tolerance  
Immunization  
 Immunological diseases  
 Mammal (Mammalia)  
 Medicine  
 Polymorphism (genetic)  
 T cell (lymphocyte)  
 Transplant (organ)  
 Transplant rejection  
 Vaccines  
 (T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Antibodies  
 TCR (T cell receptors)  
 RL: BPN (Biosynthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Class I HLA antigens  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Genes (animal)  
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (suicide; T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT Hematopoietic precursor cell  
 (tumors; T cell epitope of **minor histocompatibility**

antigen HA-1 for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

IT 204931-32-8 220419-68-1

RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study);

USES

(Uses)

(T cell epitope of **minor histocompatibility antigen HA-1** for induction of immune tolerance and for treatment of transplant rejection, graft vs. host disease, leukemia and immune disease)

RE.CNT 7

RE

(1) Den Haan, J; Eur J Immunol 1996, V26, P2680 CAPLUS

(2) Den Haan, J; Science 1995, V268, P1476 CAPLUS

(3) Den Haan, J; Science 1998, V279, P1054 CAPLUS

(4) Goulmy, E; WO 9705168 A 1997 CAPLUS

(5) Goulmy, E; WO 9705169 A 1997 CAPLUS

(6) Goulmy, E; Eye 1995, V9, P180

(7) Van Der Harst, D; Blood 1994, V83(4), P1060 CAPLUS

=> d his

(FILE 'HOME' ENTERED AT 10:38:19 ON 12 FEB 2001)

FILE 'MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS' ENTERED AT 10:38:50 ON 12 FEB 2001

L1 2960 S MINOR HISTOCOMPATIBILITY ANTIGEN

L2 19 S L1 AND DERIVATIVE

L3 13 DUP REMOVE L2 (6 DUPLICATES REMOVED)

L4 13 S L3

L5 149 S L1 AND HA-1

L6 58 DUP REMOVE L5 (91 DUPLICATES REMOVED)

L7 2 S L6 AND VLXDDLLEA

=> s 15 and VLHDDLLEA

L8 8 L5 AND VLHDDLLEA

=> dup remove 18

PROCESSING COMPLETED FOR L8

L9 2 DUP REMOVE L8 (6 DUPLICATES REMOVED)

=> d 18 1-2

L8 ANSWER 1 OF 8 MEDLINE

AN 2000166344 MEDLINE

DN 20166344

TI Molecular modeling of the **minor histocompatibility antigen HA-1** peptides binding to HLA-A alleles.

AU Ren E C; Kangueane P; Kolatkar P; Lin M T; Tseng L H; Hansen J A

CS Department of Microbiology, WHO Collaborating Center for Immunology, National University of Singapore, Singapore.. micrenec@nus.edu.sg

SO TISSUE ANTIGENS, (2000 Jan) 55 (1) 24-30.  
 Journal code: VSV. ISSN: 0001-2815.  
 CY Denmark  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200005  
 EW 20000502

L8 ANSWER 2 OF 8 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
 AN 2000049501 EMBASE  
 TI Molecular modeling of the **minor histocompatibility antigen HA-1** peptides binding to HLA-A alleles.  
 AU Ren E.C.; Kanguane P.; Kolatkar P.; Lin M.T.; Tseng L.H.; Hansen J.A.  
 CS Dr. E.C. Ren, Department of Microbiology, Faculty of Medicine, National University Singapore, Singapore 119260, Singapore. micrenec@nus.edu.sg  
 SO Tissue Antigens, (2000) 55/1 (24-30).  
 Refs: 20  
 ISSN: 0001-2815 CODEN: TSANA2

CY Denmark  
 DT Journal; Article  
 FS 022 Human Genetics  
 026 Immunology, Serology and Transplantation  
 LA English  
 SL English

=> d his

(FILE 'HOME' ENTERED AT 10:38:19 ON 12 FEB 2001)

FILE 'MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS' ENTERED AT 10:38:50 ON 12 FEB 2001

L1 2960 S MINOR HISTOCOMPATIBILITY ANTIGEN  
 L2 19 S L1 AND DERIVATIVE  
 L3 13 DUP REMOVE L2 (6 DUPLICATES REMOVED)  
 L4 13 S L3  
 L5 149 S L1 AND HA-1  
 L6 58 DUP REMOVE L5 (91 DUPLICATES REMOVED)  
 L7 2 S L6 AND VLXDDLLEA  
 L8 8 S L5 AND VLHDDLLEA  
 L9 2 DUP REMOVE L8 (6 DUPLICATES REMOVED)

=> s 15 and VLRDDLLEA

L10 5 L5 AND VLRDDLLEA

=> dup remove 110

PROCESSING COMPLETED FOR L10

L11 1 DUP REMOVE L10 (4 DUPLICATES REMOVED)

=> D 111

L11 ANSWER 1 OF 1 MEDLINE

DUPLICATE 1



AN 2000166344 MEDLINE  
 DN 20166344  
 TI Molecular modeling of the **minor histocompatibility antigen HA-1** peptides binding to HLA-A alleles.  
 AU Ren E C; Kanguane P; Kolatkar P; Lin M T; Tseng L H; Hansen J A  
 CS Department of Microbiology, WHO Collaborating Center for Immunology, National University of Singapore, Singapore.. micrenec@nus.edu.sg  
 SO TISSUE ANTIGENS, (2000 Jan) 55 (1) 24-30.  
 Journal code: VSV. ISSN: 0001-2815.  
 CY Denmark  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200005  
 EW 20000502

=> d his

(FILE 'HOME' ENTERED AT 10:38:19 ON 12 FEB 2001)

FILE 'MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS' ENTERED AT 10:38:50 ON 12 FEB 2001

L1 2960 S MINOR HISTOCOMPATIBILITY ANTIGEN  
 L2 19 S L1 AND DERIVATIVE  
 L3 13 DUP REMOVE L2 (6 DUPLICATES REMOVED)  
 L4 13 S L3  
 L5 149 S L1 AND HA-1  
 L6 58 DUP REMOVE L5 (91 DUPLICATES REMOVED)  
 L7 2 S L6 AND VLXDDLLEA  
 L8 8 S L5 AND VLHDDLLEA  
 L9 2 DUP REMOVE L8 (6 DUPLICATES REMOVED)  
 L10 5 S L5 AND VLRDDLLEA  
 L11 1 DUP REMOVE L10 (4 DUPLICATES REMOVED)

=> s l1 and HA-2

L12 55 L1 AND HA-2

=> dup remove L12

PROCESSING COMPLETED FOR L12

L13 17 DUP REMOVE L12 (38 DUPLICATES REMOVED)

=> d l13 1-17

L13 ANSWER 1 OF 17 SCISEARCH COPYRIGHT 2001 ISI (R)  
 AN 2001:76463 SCISEARCH  
 GA The Genuine Article (R) Number: 372WB  
 TI Emergence of hematopoiesis-specific **minor histocompatibility antigen** (mHag) HA-1 and HA-2 specific CD8+T cells associated with complete molecular remission after donor lymphocyte infusion (DLI) for relapsed CML.  
 AU Marijt W A F (Reprint); Kester M G D; Goulmy E; Mutis T; Drijfhout J W; Willemze R; Falkenburg J H F

CS Leiden Univ, Med Ctr, Dept Hematol, Leiden, Netherlands; Leiden Univ, Med  
Ctr, Dept Immunohematol, Leiden, Netherlands  
CYA Netherlands  
SO BLOOD, (16 NOV 2000) Vol. 96, No. 11, Part 1, pp. 478A-478A. MA 2055.  
Publisher: AMER SOC HEMATOLOGY, 1900 M STREET. NW SUITE 200, WASHINGTON,  
DC 20036 USA.  
ISSN: 0006-4971.  
DT Conference; Journal  
LA English  
REC Reference Count: 0

L13 ANSWER 2 OF 17 MEDLINE DUPLICATE 1  
AN 1999192451 MEDLINE  
DN 99192451  
TI Feasibility of immunotherapy of relapsed leukemia with ex vivo-generated  
cytotoxic T lymphocytes specific for hematopoietic system-restricted  
**minor histocompatibility antigens** [see  
comments].  
CM Comment in: Blood 1999 Dec 15;94(12):4374-6  
AU Mutis T; Verdijk R; Schrama E; Esendam B; Brand A; Goulmy E  
CS Department of Immunohematology and Blood Bank, Leiden University Medical  
Center, Leiden, The Netherlands.. Mutis@rullf2.leidenuniv.nl  
SO ~~BLOOD, (1999 Apr 1) 93 (7) 2336-41.~~  
Journal code: A8G. ISSN: 0006-4971.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
EM 199906

L13 ANSWER 3 OF 17 CAPLUS COPYRIGHT 2001 ACS  
AN 1997:215796 CAPLUS  
DN 126:198552  
TI **HA-2** antigenic peptide  
IN Goulmy, Els A. J. M.; Hunt, Donald F.; Engelhard, Victor H.  
PA Rijksuniversiteit Te Leiden, Neth.; University of Virginia Patent  
Foundation; Goulmy, Els A. J. M.; Hunt, Donald F.; Engelhard, Victor H.  
SO PCT Int. Appl., 36 pp.  
CODEN: PIXXD2

DT Patent  
LA English

FAN.CNT 1

|    | PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|----|---|------|----------|-----------------|----------|
| PI | WO 9705169  | A1   | 19970213 | WO 1996-NL183   | 19960425 |
|    | W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI |      |          |                 |          |
|    | RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN  |      |          |                 |          |
|    | US 5770201  | A    | 19980623 | US 1994-363691  | 19941223 |
|    | CA 2224909  | AA   | 19970213 | CA 1996-2224909 | 19960425 |
|    | AU 9654099  | A1   | 19970226 | AU 1996-54099   | 19960425 |
|    | AU 716907   | B2   | 20000309 |                 |          |
|    | EP 840750   | A1   | 19980513 | EP 1996-911119  | 19960425 |
|    | R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI   |      |          |                 |          |

JP 11514340 T2 19991207 JP 1996-507492 19960425  
PRAI EP 1995-202039 19950725  
WO 1996-N  
L183 19960425

L13 ANSWER 4 OF 17 CAPLUS COPYRIGHT 2001 ACS  
AN 2000:6100 CAPLUS  
DN 132:292279  
TI Nature of the **minor histocompatibility**  
**antigens**  
AU Goulmy, E.  
CS Department of Immunohaematology and Blood Bank, University Hospital,  
Leiden, 2300 RC, Neth.  
SO HLA: [Proc. Int. Histocompat. Workshop Conf.], 12th (1997), Meeting Date  
1996, Volume 2, 39-41. Editor(s): Charron, Dominique. Publisher: EDK,  
Medical and Scientific International Publisher, Sevres, Fr.  
CODEN: 68MRA5  
DT Conference; General Review  
LA English  
RE.CNT 10  
RE

(2) Den Haan, J; Eur J Immunol 1996, V26, P2680 CAPLUS

(3) Den Haan, J; Science 1995, V268, P1476 CAPLUS

(6) Goulmy, E; Curr Op Immunol 1996, V8, P75 CAPLUS

(9) Van der Harst, D; Blood 1994, V83, P1060 CAPLUS

(10) Wang, W; Science 1995, V269, P1588 CAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 17 MEDLINE DUPLICATE 2  
AN 97080610 MEDLINE  
DN 97080610  
TI Conservation of **minor histocompatibility**  
**antigens** between human and non-human primates.  
AU den Haan J M; Bontrop R E; Pool J; Sherman N; Blokland E; Engelhard V H;  
Hunt D F; Goulmy E  
CS Department of Immunohaematology and Bloodbank, Leiden University  
Hospital,  
The Netherlands.. haan.j@rulgca.leidenuniv.nl  
NC AI20963 (NIAID)  
AI33993 (NIAID)  
SO EUROPEAN JOURNAL OF IMMUNOLOGY, (1996 Nov) 26 (11) 2680-5.  
Journal code: EN5. ISSN: 0014-2980.  
CY GERMANY: Germany, Federal Republic of  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Cancer Journals; Priority Journals  
EM 199703

L13 ANSWER 6 OF 17 BIOSIS COPYRIGHT 2001 BIOSIS  
AN 1996:111037 BIOSIS  
DN PREV199698683172  
TI Mismatches of **minor histocompatibility**  
**antigens** between HLA-identical donors and recipients and the  
development of graft-versus-host disease after bone marrow  
transplantation.  
AU Goulmy, Els (1); Schipper, Ronald; Pool, Jos; Blokland, Els; Falkenburg,  
J. H. Frederick; Vossen, Jaak; Gratwohl, Alois; Vogelsang, Georgia B.;  
Van

Houwelingen, Hans C.; Van Rood, Jon J.  
CS (1) Dep. Immunohematology Blood Bank, Leiden Univ. Hosp., P.O. Box 9600,  
2300 RC Leiden Netherlands  
SO New England Journal of Medicine, (1996) Vol. 334, No. 5, pp. 281-285.  
ISSN: 0028-4793.  
DT Article  
LA English

L13 ANSWER 7 OF 17 CAPLUS COPYRIGHT 2001 ACS  
AN 1996:532658 CAPLUS  
DN 125:193403  
TI Functional expression of **minor histocompatibility**  
**antigens** on human peripheral blood dendritic cells and epidermal  
Langerhans cells  
AU Van Lochem, Ellen; Van Der Keur, Maarten; Mommaas, A. Mieke; De Gast,  
Gijsbert C.; Goulmy, Els  
CS Department Immunohematology and Bloodbank, Leiden University Hospital,  
Leiden, 2300 RC, Neth.  
SO Transplant Immunol. (1996), 4(2), 151-157  
CODEN: TRIME2; ISSN: 0966-3274  
DT Journal  
LA English

L13 ANSWER 8 OF 17 MEDLINE DUPLICATE 3  
AN 95288637 MEDLINE  
DN 95288637  
TI Identification of a graft versus host disease-associated human  
**minor histocompatibility antigen**.  
AU den Haan J M; Sherman N E; Blokland E; Huczko E; Koning F; Drijfhout J W;  
Skipper J; Shabanowitz J; Hunt D F; Engelhard V H; et al  
CS Department of Immunohaematology, University Hospital, Leiden,  
Netherlands.  
NC AI33993 (NIAID)  
AI20963 (NIAID)  
SO SCIENCE, (1995 Jun 9) 268 (5216) 1476-80.  
Journal code: UJ7. ISSN: 0036-8075.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199509

L13 ANSWER 9 OF 17 MEDLINE DUPLICATE 4  
AN 95362850 MEDLINE  
DN 95362850  
TI Recognition of clonogenic leukemic cells, remission bone marrow and  
HLA-identical donor bone marrow by CD8+ or CD4+ **minor**  
**histocompatibility antigen**-specific cytotoxic T  
lymphocytes.  
AU Faber L M; van der Hoeven J; Goulmy E; Hooftman-den Otter A L; van  
Luxemburg-Heijs S A; Willemze R; Falkenburg J H  
CS Department of Hematology, University Medical Center, Leiden, The  
Netherlands..  
SO JOURNAL OF CLINICAL INVESTIGATION, (1995 Aug) 96 (2) 877-83.  
Journal code: HS7. ISSN: 0021-9738.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
EM 199511

L13 ANSWER 10 OF 17 MEDLINE DUPLICATE 5

AN 94333484 MEDLINE

DN 94333484

TI Presentation of viral antigens restricted by H-2Kb, Db or Kd in proteasome

subunit LMP2- and LMP7-deficient cells.

AU Zhou X; Momburg F; Liu T; Abdel Motal U M; Jondal M; Hammerling G J; Ljunggren H G

CS Microbiology and Tumor Biology Center, Karolinska Institute, Stockholm, Sweden.

SO EUROPEAN JOURNAL OF IMMUNOLOGY, (1994 Aug) 24 (8) 1863-8.

Journal code: EN5. ISSN: 0014-2980.

CY GERMANY: Germany, Federal Republic of

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; Cancer Journals

EM 199411

L13 ANSWER 11 OF 17 MEDLINE DUPLICATE 6

AN 94154267 MEDLINE

DN 94154267

TI Recognition of **minor histocompatibility antigens** on lymphocytic and myeloid leukemic cells by cytotoxic T-cell clones.

AU van der Harst D; Goulmy E; Falkenburg J H; Kooij-Winkelaar Y M; van Luxemburg-Heijs S A; Goselink H M; Brand A

CS Department of Immunohematology and Bloodbank, University Medical Center, Leiden, The Netherlands..

SO BLOOD, (1994 Feb 15) 83 (4) 1060-6.

Journal code: A8G. ISSN: 0006-4971.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals

EM 199406

L13 ANSWER 12 OF 17 MEDLINE DUPLICATE 7

AN 94083660 MEDLINE

DN 94083660

TI **Minor histocompatibility antigens** HA-1-, -2-, and -4-, and HY-specific cytotoxic T-cell clones inhibit human hematopoietic progenitor cell growth by a mechanism that is dependent on direct cell-cell contact.

AU Marijt W A; Veenhof W F; Goulmy E; Willemze R; van Rood J J; Falkenburg J H

CS Department of Hematology, University Medical Center, Leiden, The Netherlands..

SO BLOOD, (1993 Dec 15) 82 (12) 3778-85.

Journal code: A8G. ISSN: 0006-4971.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals

EM 199403

L13 ANSWER 13 OF 17 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.DUPLICATE 8  
 AN 93095092 EMBASE  
 DN 1993095092  
 TI Isolation of an HLA-A2.1 extracted human minor histocompatibility peptide.  
 AU De Bueger M.; Verreck F.; Blokland E.; Drijfhout J.W.; Amous R.; Koning F.; Goulmy E.  
 CS Department of Immunohaematology, University Hospital Leiden, Rijnsburgerweg 10,NL-2333 AA Leiden, Netherlands  
 SO European Journal of Immunology, (1993) 23/3 (614-618).  
 ISSN: 0014-2980 CODEN: EJIMAF  
 CY Germany  
 DT Journal; Article  
 FS 026 Immunology, Serology and Transplantation  
 029 Clinical Biochemistry  
 LA English  
 SL English

L13 ANSWER 14 OF 17 MEDLINE DUPLICATE 9  
 AN 93246305 MEDLINE  
 DN 93246305  
 TI A genetic analysis of human **minor histocompatibility antigens** demonstrates Mendelian segregation independent of HLA.

AU Schreuder G M; Pool J; Blokland E; van Els C; Bakker A; van Rood J J; Goulmy E  
 CS Department of Immunohaematology, University Hospital Leiden, The Netherlands..  
 SO IMMUNOGENETICS, (1993) 38 (2) 98-105.  
 Journal code: GI4. ISSN: 0093-7711.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199308

L13 ANSWER 15 OF 17 MEDLINE DUPLICATE 10  
 AN 92373026 MEDLINE  
 DN 92373026  
 TI Tissue distribution of human **minor histocompatibility antigens**. Ubiquitous versus restricted tissue distribution indicates heterogeneity among human cytotoxic T lymphocyte-defined non-MHC antigens.

AU de Bueger M; Bakker A; Van Rood J J; Van der Woude F; Goulmy E  
 CS Department of Immunohaematology, University Hospital, Leiden, The Netherlands..  
 SO JOURNAL OF IMMUNOLOGY, (1992 Sep 1) 149 (5) 1788-94.  
 Journal code: IFB. ISSN: 0022-1767.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
 EM 199211

L13 ANSWER 16 OF 17 CAPLUS COPYRIGHT 2001 ACS  
 AN 1992:233346 CAPLUS  
 DN 116:233346  
 TI Transfected human class I gene product adequately assembles **minor**

**histocompatibility antigens**  
 AU Goulmy, Els; Pool, Jos; Blokland, Els; Geraghty, Dan  
 CS Dep. Immunohaematol., Univ. Hosp., Leiden, 2300 RC, Neth.  
 SO Immunogenetics (1991), 34(4), 270-2  
 CODEN: IMNGBK; ISSN: 0093-7711  
 DT Journal  
 LA English

L13 ANSWER 17 OF 17 MEDLINE DUPLICATE 11  
 AN 89067836 MEDLINE  
 DN 89067836  
 TI Cellularly defined **minor histocompatibility**  
**antigens** are differentially expressed on human hematopoietic  
 progenitor cells.  
 AU Voogt P J; Goulmy E; Veenhof W F; Hamilton M; Fibbe W E; Van Rood J J;  
 Falkenburg J H  
 CS Department of Hematology, University Medical Center, Leiden, The  
 Netherlands..  
 SO JOURNAL OF EXPERIMENTAL MEDICINE, (1988 Dec 1) 168 (6) 2337-47.  
 Journal code: I2V. ISSN: 0022-1007.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English

---

FS Priority Journals; Cancer Journals  
 EM 198903

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

|  |            |         |
|--|------------|---------|
| COST IN U.S. DOLLARS                       | SINCE FILE | TOTAL   |
|  | ENTRY      | SESSION |
| FULL ESTIMATED COST                        | 66.79      | 66.94   |
| DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) | SINCE FILE | TOTAL   |
|  | ENTRY      | SESSION |
| CA SUBSCRIBER PRICE                        | -4.12      | -4.12   |

STN INTERNATIONAL LOGOFF AT 10:50:41 ON 12 FEB 2001